AJMS
Asian Journal of Mathematical Sciences

RESEARCH ARTICLE

SOLUTION OF BEAL'S CONJECTURE IN THE PARADIGM OF QUANTUM MATHEMATICS (AN INTRODUCTION INTO THE REALM OF QUANTUM MATHEMATICS)

*** YURI A. IVLIEV**

Independent researcher
Dr. of Science, full member of the International Informatization Academy Moscow, Russia.

Corresponding Email: yuri.ivliev@gmail.com

Received: 17-07-2026; Revised: 19-07-2026; Accepted: 05-09-2026

ABSTRACT

The article gives a tight review of the author's work dedicated to a new trend in modern mathematics called quantum computing mathematics. Quantum computing methods are interpreted extendedly as relating to quantum mathematical information (quantum mathematics). Mathematical information quanta are whole numbers (integers) characterizing principal integrity of relevant mathematical objects. This approach was applied to solving Beal's Conjecture (Generalized Fermat's Last Theorem) that allowed not only to prove the famous problem but unmask excessive formalism of traditional quantum theories. It was revealed also that Fermat's method of infinite descent was in fact the first case of exact application of quantum ideology in pure mathematics.

Keywords: Beal's Conjecture, Fermat's Last Theorem, arithmetic geometry, partitions, quantum mathematics.,

INTRODUCTION

Beal's Conjecture [1-2] deals with arbitrary positive whole powers of natural numbers except the second combined in one equation similar to the well-known equation of Fermat's Last Theorem. Among all well-known mathematics conjectures Beal's Conjecture is occupying a peculiar place being a formal generalization of Fermat's Last Theorem The Beal proposition can be solved by the ancient Greek arithmetic geometry methods applied successfully as well to the Fermat problem and expanded with Fermat's method of infinite descent. Pierre de Fermat formulated his proposition on the margin of Diophantus's Arithmetic (near the task 8 of the book II). The eighth problem of the second book asks to separate a square into two squares in whole numbers. It was known long ago that this problem has an infinite set of solutions. But Fermat generalized the task in case of any whole power above the second and pointed out at impossibility of such partition in whole numbers claiming here that he found a "miraculous"

proof of this proposition. In order to reconstruct Fermat's proof it is necessary to understand what Fermat meant by his recording on the margins of Diophantus's book. So at the very beginning it is a question of fundamental approach to this problem-solving by Fermat. Psychologically we can assume that Fermat applied a phenomenological approach when each power of natural numbers was considered as an amount of indivisible multidimensional unit «cubes» (squares) in multidimensional arithmetic space. Thus the task was analogous to the ancient receipt of alchemy: first resolve the power into elementary units and then put them together in a required manner. "Pure" mathematical units can be chosen as such elementary units but unlike the quantum bit they can have the structure of multidimensional arithmetic space in the case of whole powers.

How could Fermat solve this unique problem straight off and without a shadow of doubt ? The sole reason for it is that he saw the mental picture of his proof. Such a picture emerged in his consciousness during his insight allowing him to investigate instantly all necessary details of solution [1-2]. Visual image of the problem must have had a geometrical form, which apparently could not take its place on narrow margins. This geometric pattern serves as general illustration for Euclid's theorem about mean proportionals, from which formulation of Pythagorean theorem and Fermat's proposition (called Fermat's Last Theorem later on) could be easily derived. Fig. 1 taken from [1-2] shows stylized design of Euclid's geometrical theorem on the fractal surface of similar right angle triangles at an instantaneous position of the small diameter of Fig. 1 shifting from state Φ1 to state Φ2.




Fig. 1 The image of Beal's Conjecture mathematical quantum (the designations are explained in the text)

Mental perception of n-th degrees of whole powers can be related to Cartesian product of whole numbers from n-dimensional arithmetic space and then each n-th power could be represented as a collection of n-dimensional unit cubes transferred from the state Φ1 to the state Φ2 one by one on the diagram of Fig. 1. This procedure serves as well for finding Pythagorean triples by the method of sorting square units one by one in Pythagorean equation. This method can be applied also to the equation of generalized Fermat's Last Theorem in order to show that splitting of n-th power of whole numbers into two other powers with $n > 2$ is impossible. As it will be seen further, the bases of whole powers in Beal's Conjecture can consist only of Pythagorean triples in the form of Pythagorean partitions in order to satisfy quantum splitting of whole powers into other ones.

Solution of Beal's Conjecture and Fermat's Last Theorem.

Let us write hypothetical Beal's Conjecture equality [1] in the following way:

$$z^n = x^n + y^n \qquad (1)$$

with positive integers x, y, z having a common factor and exponent n taking simultaneously the next spectrum of values: n = (m, p, q) , where integers p, q, m at least 3 and n has one independent value for each term. So we suppose at the beginning that equality (1) exists and partitions of the type (1) can be obtained. This method of proving is related to plausible reasoning and called the rule of contraries.

Consider equality (1) as a partition of whole number zn into two non-zero whole parts xn and yn consisting of whole numbers. Equality (1) resembles Pythagorean equation in real numbers, if we could bring powers in (1) to the degree 2 with whole parts in the similar partition: z2 = xn/ zn–2 + yn/ zn–2. To produce such scaling, the notion of right-angled numbers was introduced (these numbers are different from so called right angle triangle numbers representing Pythagorean triples).

Definition.

Right-angled number is such a non-negative real number, the square of which is a whole non-negative number.

The set of right-angled numbers P = {0, 1, √2 , √3 , 2, √5, …} is countable. The system of right-angled numbers P = □P,+,·,0,1□ is defined by operations of addition and multiplication and two singled out elements (zero and unit). The system P is non-closed in relation to addition. Notice that the set of non-negative whole numbers is a subset of the set of right-angled numbers. Then consider (1) on the 2-dimensional lattice of right-angled numbers with coordinates xo , yo, and that, which we call the norms of a right-angled number z assigned to different pairs (xo , yo) and differing from each other by the value of its summands: z2 = xo2 + yo2. The norm of non-zero right-angled numbers is always whole and cannot be less than 1.

For the purpose of reducing (1) to the view of Pythagorean equation in the system of right-angled numbers, one can rewrite (1) as an equality for some coprime x', y', z', and common whole factor d: (x'd)p + (y'd)q = (z'd)m and fulfil scaling-down:

$$(z'd)^2 = (x'd)^p / (z'd)^{m-2} + (y'd)^q / (z'd)^{m-2} = (x')^p\, d^{\,p-m+2} / (z')^{m-2} + (y')^q\, d^{\,q-m+2} / (z')^{m-2} = x_o^{\,2} +$$

yo2 , where xo2 and yo2 with appropriate d are squares of some right-angled numbers xo and yo. Right-angled numbers correspond to the wave functions of traditional quantum theories

They assume the following view of (1) after fulfilling scaling-up:

$$z^m = x^p + y^q = z^{m-2}\,(x_o^{\,2} + y_o^{\,2}) \qquad (2)$$

with whole right-angled numbers xo and yo , if p,q = m [8], .presenting the first Pythagorean triple in the series of the next scaling-down of less Pythagorean equations.

Let us apply now the ancient method of making powers using Euclid's geometrical theorem and produce two chains of proportions connected with each other with some equality

presenting integer z as a sum of two whole numbers:

$$z/x_o = x_o/k = k/k_1 = \ldots = k_{m-3}/k_{m-2} \qquad (3)$$

$$z/y_o = y_o/l = l/l_1 = \ldots = l_{m-3}/l_{m-2}$$

where z, xo , yo are right-angled numbers from (2), m natural index at least 3,and z=k + l ; k and l are some whole parts of z taken from the method of scaling-down.

From proportions (2) one can obtain the next formulae

$$x_o^2 = kz = (k_1 z / x_o)z\ ,\ x_o^3 = k_1 z^2 = (k_2 z / x_o)z^2\ , \ldots ,\ x_o^m = k_{m-2}\, z^{m-1}, \qquad (4)$$

$$y_o^2 = lz = (l_1 z / y_o)z\ ,\quad y_o^3 = l_1 z^2 = (l_2 z / y_o)z^2\ , \ldots ,\ y_o^m = l_{m-2}\, z^{m-1},$$

where integers k and l are found from the basic equality (1):

$$z = (z'd) = (x'd)^p / (z'd)^{m-1} + (y'd)^q / (z'd)^{m-1} = (x')^p\, d^{\,p-m+1} / (z')^{m-1} + (y')^q\, d^{\,q-m+1} / (z')^{m-1} = k + l$$

If exponents p and q more or equal m, then numbers k and l are whole with d = (z')m–1 as a minimum (d can be some whole number divisible by this minimum).

From (2) we get equal similar partitions of zn into two whole parts:

$$z^m = x^p + y^q = z^{m-2}(x_o^2 + y_o^2) = x^m + y^m, \qquad (5)$$

where xp = (xp/m)m = xm, yq = (yq/m)m = ym with whole x, y by construction (for simplicity we do not change here the designations for x, y although exponents p and q can differ from m, moreover irrational numbers xp/m and yq/m do not fit in Procrustean bed of quantum paradigm). Square roots of xm, ym are mean proportionals between xo2 and zm–2, yo2 and zm–2 describing a bigger right angle triangle defined by the hidden Pythagorean equality zm = xm + ym found from the relations: xm = k zm–1, ym = l zm–1. This implicit triangle is similar to that with sides z , xo , yo represented by equality z2 = xo2 + yo2.

So (1) comes to the Fermat equality (Pythagorean equality in natural numbers) that is equivalent to hypothetical phenomenological equality (1);

$$z^m = x^m + y^m,\ m \geq 3 \qquad (6)$$

with whole x = x'd, y = y'd, z = z'd, and some whole factor d that can be expanded into the product of prime factors. One can now prove Fermat's Last Theorem using the same methods as above in order to obtain solution of the Beal Conjecture in full and one measure.

Let us write Fermat's Last Theorem in its usual form of hypothetical equation:

$$z^n = x^n + y^n\ ,\ n > 2\ ,\quad x,\ y, z \text{ integers}. \qquad (7)$$

Suppose that one solution at least was found. Then we shall try to construct such a solution and make certain of its impossibility. We shall work in the system of right-angled numbers.

To get powers of whole numbers presented in (7), let us produce two chains of continued proportions connected with each other by the norm z2 = x02 + y02 :

$$z/x_0 = x_0/k = k/k_1 = \ldots = k_{n-3}/k_{n-2}$$

$$z/y_0 = y_0/l = l/l_1 = \ldots = l_{n-3}/l_{n-2} \qquad (8)$$

where natural indices of the last terms of each chain in (8) are obtained from n > 2. Continued proportions (8) yield the following formulae:

$$kz = x_0^2,\ k_1 z = x_0 k,\ k_2 z = x_0 k_1,\ \ldots,\ k_{n-2} z = x_0 k_{n-3} \qquad (9)$$

$$lz = y_0^2,\ l_1 z = y_0 l,\ l_2 z = y_0 l_1,\ \ldots,\ l_{n-2} z = y_0 l_{n-3}$$

$$x_0^2 = kz = (k_1 z / x_0) z,\quad x_0^3 = k_1 z^2 = (k_2 z / x_0) z^2,\ \ldots,\ x_0^n = k_{n-2} z^{n-1} \tag{10}$$

$$y_0^2 = lz = (l_1 z / y_0) z,\quad y_0^3 = l_1 z^2 = (l_2 z / y_0) z^2,\ \ldots,\ y_0^n = l_{n-2} z^{n-1}$$

It is necessary to fix now the norm for the partition zn into two like powers in (7). As in the case of Beal's Conjecture, let us assume that z, x, y in presupposed equality (7) have a common factor d :

$z = (z'd),\ x = (x'd),\ y = (y'd)$, where $z',\ x'.\ y'$ coprime. Thereupon divide equality (7) by z^{n-1} and get: $z = (z'd) = (x'd)^n / (z'd)^{n-1} + (y'd)^n / (z'd)^{n-1} = k + l$, where k and l integers with $d = (z')^{n-1}$ as a minimum. From this and (9)-(10) it follows that $z^2 = x_0^2 + y_0^2$ and $z^n = z^{n-2} (x_0^2 + y_0^2)$ is a scaled-up modification of the norm $z^2 = x_0^2 + y_0^2$.

Further, one can obtain a singular partition of zn into three terms from (10) for the given norm when n > 2 :

$$z^n = x_0^n + y_0^n + \lambda_n \tag{11}$$

where $\lambda n = z^{n-1} [(k - k_{n-2}) + (l - l_{n-2})]$ is a remainder after subtracting x_0^n and y_0^n out of z^n such that $\lambda_n > 0$ when $n > 2$ and $x_0 y_0 \neq 0$, $\lambda_n = 0$ when $n = 2$ and $x_0 y_0 \neq 0$, $x_0,\ y_0 \in [0, z]$, $z \in (0, \infty)$.

Thus there exists one-to-one correspondence between each pair of numbers (x0 , y0) with norm z2 = x02 + y02 from 2-dimensional arithmetic space and each corresponding partition of any whole power with n > 2 of integer z from n-dimensional arithmetic space into the sum of the same powers of numbers x0 , y0 and remainder λn from (11). Isomorphism (one-to-one correspondence) between the set of points of 2-dimensional Euclidean space with position vector length z and coordinates x0 , y0 , the set of partitions of z2 into squares, and the sets of partitions (11) for any whole n > 2 can be written as follows:

$$\left\{ z \Rightarrow (xo\ , yo) \right\} \leftrightarrow \{ z^2 = x_0^2 + y_0^2 \} \leftrightarrow \{ z^n = x_0^n + y_0^n + \lambda_n \},$$

where sets of partitions are generated by the next power similarities:

$$z \leftrightarrow z^2 \leftrightarrow z^n ,\ x_0 \leftrightarrow x_0^2 \leftrightarrow x_0^n ,\ y_0 \leftrightarrow y_0^2 \leftrightarrow y_0^n .$$

Partitions (11) can be reduced to the norm, from which they were obtained:

$$z^n = x_0^n + y_0^n + \lambda_n = z^{n-2} (x_0^2 + y_0^2) = x^n + y^n \tag{12}$$

Formula (12) represents by itself a combinatorial equality of two partitions in three and two terms because of the one-to-one correspondence between pairs (x0 , y0) and presupposed partition (7). It means that partition (11) coincides with partition (7) if the latter exists. In the case of right-angled numbers this equality is realized only if x0, y0 integers [8]. Algorithm of such correspondence is given in the next formula (13). Thus scaling invariance of the norm z2 = (x02 + y02) leads to the following equalities of different fragments of partitions (12):

$$x_0^n + y_0^n = (x^n \text{ or } y^n) \tag{13}$$

and correspondingly λn = (yn or xn). It can be noticed that x0n ≠ zn-2·y02= yn and y0n ≠ zn-2·x02 = xn because of the lack of coincidence of decompositions in factorization of numbers x0n and yn, y0n and xn

. Obviously, $x0n \neq zn\text{-}2 \cdot x02$ and $y0n \neq zn\text{-}2 \cdot y02$. One can show also that $x0$ and $y0$ cannot be irrational in (13) on account of integer partition of zn into xn and yn when $n > 2$.

Let us come back to the assumption at the beginning of the proof that integer solution (7) exists. This assumption is substantiated only if there is a concrete solution (13) in whole numbers. In order to check the validity of (13) it is necessary to construct it with the same reasoning as before, since equations (7) and (13) are identical by their properties. This procedure can be continued to infinity in the direction of decreasing whole numbers under condition that sequence of different chained equalities never stops and numbers x02 and y02 in (12) will be always whole. If it is not so, i.e., x02 and y02 in chained equalities (13) turn out to be fractions, then this means that solution (7) does not exist in the system of right-angled numbers (the base z can be splitted only into whole numbers by assumption). On the other hand, infinite sequence of chained equalities (13) leads to infinite decreasing of positive whole numbers that is impossible and therefore assuming that there exists an integer solution of (7) , when $n > 2$, is not true. Thus the theorem is proved both for all even and for all odd degrees of whole numbers and for any finite whole x, y, z, d.

CONCLUSION

Beal's Conjecture solution contains in itself the description of a new hypothetical mathematical object with simple properties conditioned only by its intrinsic structure. This mathematical object represents by itself a quantum of mathematical information displayed on Fig. 1 and pictured as dichotomic vector having two independent states Φ1 and Φ2 in 2-dimensional information space. Being applied to Beal's Conjecture, these states can be written as follows: $\Phi 2 = | zn + 0 >$, $\Phi 1 = | 0 + zn >$, Φ (superposition of the states Φ1 and Φ2) $= | xn + yn >$., The dichotomic vector symbolized by a small diameter turns as if inside out in 1-dimensional space of human visual perception, but in fact it carries out two independent rotations in 2-dimentional information space of human mental perception preserving its orientation in the range of values from 0 to 1.

Conception of quantum information can be related not only to specific mathematical objects but to the content of other sciences using mathematical items and methods in research of their specific quantum objects and their transfer. Thus quantum mathematics integrates all other relevant sciences (quantum physics, quantum chemistry, quantum biology, quantum psychology, etc.) from the point of view of unified quantum information space. The doubling of dimension of quantum information transfer opens the way to discover new possibilities of motion and transformation in inner quantum space resulting also in outer space of perception and producing such unusual effects as levitation, teleportation, and other super-phenomena. Perhaps super-precise technologies of the ancients were based on technical application of quantum information principles being the main secret of highly developed lost civilizations.

This review of quantum solution of Beal's Conjecture rests on the author's previous works having done essential contribution to this problem-solving [3-9]. It was posted first in [10] with some inaccuracies relating to Fig.1 and states Φ1 , Φ2 distorting the true image of them but not vitiating the following calculations of quantum equalities leading to the end of the proof.

REFERENCES

1. Ivliev, Y. A., Phenomenological approach to solution of Beal's Conjecture and Fermat's Last Theorem. European Journal of Natural History 2017 № 4, 54-59.
2. Ivliev, Y. A., Beal's Conjecture as global breakthrough in natural sciences. International Conference "Global Science and Innovation" December 17-18, 2013, Materials of the I International Scientific Conference, Vol. II, USA, Chicago 2013, , 345-349.
3. Ivliev, Y. A., On natural units of information interaction in the universe. Engineering and Automation Problems 1996 № 5-6, 13-17.
4. Ivliev, Y. A., On instanton anatomy. Superconductivity: Research and Development 1996 № 7-8, 23-29.
5. Ivliev, Y. A., From mechanics after Newton and Einstein to generalized relativity mechanics. Engineering and Automation Problems 1998 № 4, 88-90.
6. Ivliev, Y. A., Quantum and pseudoquantum teleportation as effects of generalized relativity mechanics. Engineering and Automation Problems 2000 Vol. 1 Special issue № 1, 68-71.
7. Ivliev, Y. A., Fermat's Last Theorem is a quantum theorem in the realm of quantum information. International Journal of Applied and Fundamental Research 2010 № 2, 17-20.
8. Ivliev , Y.A., Reconstruction of nativus proof of Fermat's Last Theorem. The Integrated Scientific Journal 2006 № 7, 3 - 9.
9. URL: http: //yuri-andreevich-ivliev.narod.ru/
10. Ivliev , Y.A., Solution of Beal's Conjecture in the Paradigm of Quantum Mathematics. Asian Journal of Mathematical Sciences. Jan.- Feb. 2018 vol.2 issue 1, 1-5.